

# Involvement of N-6 Adenine-Specific DNA Methyltransferase 1 (*N6AMT1*) in Arsenic Biomethylation and Its Role in Arsenic-Induced Toxicity

Xuefeng Ren,<sup>1\*</sup> Maria Aleshin,<sup>1</sup> William J. Jo,<sup>2</sup> Russel Dills,<sup>3</sup> David A. Kalman,<sup>3</sup> Christopher D. Vulpe,<sup>2</sup> Martyn T. Smith,<sup>1</sup> and Luoping Zhang<sup>1</sup>

<sup>1</sup>Genes and Environment Laboratory, Division of Environmental Health Sciences, School of Public Health, and <sup>2</sup>Department of Nutritional Sciences and Toxicology, University of California–Berkeley, Berkeley, California, USA; <sup>3</sup>Department of Environmental and Occupational Health Sciences, School of Public Health and Community Medicine, University of Washington, Seattle, Washington, USA

**BACKGROUND:** In humans, inorganic arsenic (iAs) is metabolized to methylated arsenical species in a multistep process mainly mediated by arsenic (+3 oxidation state) methyltransferase (AS3MT). Among these metabolites is monomethylarsonous acid (MMA<sup>III</sup>), the most toxic arsenical species. A recent study in *As3mt*-knockout mice suggests that unidentified methyltransferases could be involved in alternative iAs methylation pathways. We found that yeast deletion mutants lacking *MTQ2* were highly resistant to iAs exposure. The human ortholog of the yeast *MTQ2* is N-6 adenine-specific DNA methyltransferase 1 (*N6AMT1*), encoding a putative methyltransferase.

**OBJECTIVE:** We investigated the potential role of *N6AMT1* in arsenic-induced toxicity.

**METHODS:** We measured and compared the cytotoxicity induced by arsenicals and their metabolic profiles using inductively coupled plasma–mass spectrometry in UROtsa human urothelial cells with enhanced *N6AMT1* expression and UROtsa vector control cells treated with different concentrations of either iAs<sup>III</sup> or MMA<sup>III</sup>.


**RESULTS:** *N6AMT1* was able to convert MMA<sup>III</sup> to the less toxic dimethylarsonic acid (DMA) when overexpressed in UROtsa cells. The enhanced expression of *N6AMT1* in UROtsa cells decreased cytotoxicity of both iAs<sup>III</sup> and MMA<sup>III</sup>. Moreover, *N6AMT1* is expressed in many human tissues at variable levels, although at levels lower than those of *AS3MT*, supporting a potential participation in arsenic metabolism *in vivo*.

**CONCLUSIONS:** Considering that MMA<sup>III</sup> is the most toxic arsenical, our data suggest that *N6AMT1* has a significant role in determining susceptibility to arsenic toxicity and carcinogenicity because of its specific activity in methylating MMA<sup>III</sup> to DMA and other unknown mechanisms.

**KEY WORDS:** arsenic methylation, arsenic toxicity, arsenite, monomethylarsonous acid, *N6AMT1*. *Environ Health Perspect* 119:771–777 (2011). doi:10.1289/ehp.1002733 [Online 30 December 2010]

Inorganic arsenic (iAs) compounds are considered known human carcinogens that target multiple sites, including the lung, skin, and urinary bladder [International Agency for Research on Cancer (IARC) 2004; Pershagen 1981; Smith et al. 1992; Smith and Steinmaus 2009; Straif et al. 2009]. In addition, chronic exposure to high levels of iAs has been associated with the development of multiple diseases and deleterious health effects in humans (Abernathy et al. 1999; Kapaj et al. 2006).

In humans, as in many animal species, iAs is metabolized to monomethylarsonous acid (MMA) and dimethylarsonic acid (DMA). The most cited conceptual model of arsenic methylation involves the reduction of pentavalent iAs (iAs<sup>V</sup>) to trivalent iAs (iAs<sup>III</sup>), with subsequent methylation (Drobna et al. 2009). The general scheme is as follows:



Among these metabolites, MMA<sup>III</sup> is the most toxic arsenical species (Drobna et al. 2005; Ferrario et al. 2008; Kligerman et al. 2003; Petrick et al. 2001). It is generally accepted

that arsenic (+3 oxidation state) methyltransferase (AS3MT) is responsible for catalyzing methyl group transfer from *S*-adenosyl methionine (SAM) to iAs (Thomas et al. 2007). However, a recent study by Drobna et al. (2009) showed that knockout of *As3mt* in the mouse does not completely abolish the methylation of iAs, suggesting that there are alternative pathways for arsenic methylation in these animals. Although Bentley and Chasteen (2002) and Hall et al. (1997) suggested that arsenic methylation could be due to gastrointestinal tract microbiota, they also speculated that unidentified methyltransferases may be responsible for the methylated arsenicals found in *As3mt*-knockout mice.

We conducted a genomewide, parallel phenotypic screen of yeast deletion mutants to identify the genes required for the growth of yeast in the presence of MMA<sup>III</sup> and iAs<sup>III</sup> (Jo et al. 2009). We found a yeast strain with deletion of *MTQ2*, which was highly resistant to iAs<sup>III</sup>. *MTQ2* encodes a SAM-dependent methyltransferase and has been shown to be involved in the methylation of release factor eRF1 in yeast (Polevoda et al. 2006). The human ortholog of the yeast *MTQ2* is N-6 adenine-specific DNA methyltransferase 1 (*N6AMT1*), a putative methyltransferase. Although the bacterial homologs

of *N6AMT1* have been shown to methylate DNA N6-adenine (Stephens et al. 1996), the current data do not indicate its function in the methylation of adenine in the DNA of mammalian cells (Ratel et al. 2006).

Our goal in the present study was to explore the mechanism by which *N6AMT1* confers resistance to arsenic toxicity. We enhanced *N6AMT1* gene expression in UROtsa cells, given its relatively low expression in these cells. The UROtsa cell line, originally isolated from a primary culture of normal human uroepithelium, does not methylate arsenic because of the absence of *AS3MT* expression (Drobna et al. 2005; Styblo et al. 2000) and has been used as a model for bladder epithelium and arsenic-induced bladder cancer (Bredfeldt et al. 2006; Eblin et al. 2008; Sens et al. 2004). Here, we show that *N6AMT1* is a human methyltransferase specifically involved in the biomethylation of MMA<sup>III</sup> to DMA. Given that MMA<sup>III</sup> is the most toxic arsenical and its implication in arsenic toxicity and carcinogenicity, *N6AMT1* may have a significant role in modulating arsenic-induced toxicity and carcinogenicity.

## Materials and Methods

**Cultures of yeast strains and human UROtsa cells.** The wild-type BY4743 yeast strain was

Address correspondence to L. Zhang, School of Public Health, University of California–Berkeley, B84 Hildebrand Hall MC #7356, Berkeley, CA 94720 USA. Telephone: (510) 643-5189. Fax: (510) 642-0427. E-mail: luoping@berkeley.edu

Supplemental Material is available online (doi:10.1289/ehp.1002733 via <http://dx.doi.org/>).

\*Current address: Department of Social and Preventive Medicine, School of Public Health and Health Professions, State University of New York at Buffalo, Buffalo, NY 14214 USA.

We thank J. Gandolfi (University of Arizona–Tucson) for the generous gift of MMA<sup>III</sup> iodide and P. Simeonova (National Institute for Occupational Safety and Health) for providing UROtsa cells. We also thank C. McHale and N. Dangleben for assistance with manuscript preparation.

This research was funded by the Superfund Research Program through National Institute of Environmental Health Sciences grant P42 ES004705 (to M.T.S., C.D.V., and L.Z.).

The authors declare they have no actual or potential competing financial interests.

Received 16 July 2010; accepted 29 December 2010.

purchased from Invitrogen (Carlsbad, CA), and the *MTQ2* deletion strain has the same background as the wild-type strain. Growth was conducted in rich media [yeast extract-peptone-dextrose (YPD)] at 30°C with shaking at 200 rpm. UROtsa cells (generously provided by P. Simeonova, National Institute for Occupational Safety and Health, Morgantown, WV) were cultured at a starting cell density of  $4\text{--}5 \times 10^4$  cells/mL in RPMI 1640 (Mediatech, Inc., Manassas, VA) with L-glutamine, 10% fetal bovine serum, 100 IU/mL penicillin, and 100 µg/mL streptomycin (Omega Scientific, San Diego, CA), under standard culturing conditions.

**Arsenical exposures.** We purchased sodium arsenite [ $\text{NaAsO}_2$  ( $\text{iAs}^{\text{III}}$ ); purity > 99%] from Sigma-Aldrich (St. Louis, MO). Diiodomethylarsine [ $\text{MMA}^{\text{III}}$  iodide ( $\text{MMA}^{\text{III}}$ )] was a generous gift from J. Gandolfi (University of Arizona, Tucson, AZ).  $\text{iAs}^{\text{III}}$  and  $\text{MMA}^{\text{III}}$  solutions were freshly prepared using sterile water (Milli-Q; Millipore, Billerica, MA) and protected from light before use. Yeast cells were treated with either  $\text{iAs}^{\text{III}}$  or  $\text{MMA}^{\text{III}}$  at concentrations ranging from 0 to 300 µM. Once UROtsa cells reached 70–80% confluence in culture, they were treated with  $\text{iAs}^{\text{III}}$  at concentrations from 0 to 100 µM or  $\text{MMA}^{\text{III}}$  at 0–5 µM.

**Yeast growth assay.** Yeast strains were pregrown in YPD media to mid-log phase, diluted in fresh media to an optical density at 595 nm ( $\text{OD}_{595}$ ) of 0.0165, and inoculated into a 48-well microplate. Stock solutions of arsenicals were added to each culture with at least three replicate wells per dose.

Plates were incubated in a Tecan GENios spectrophotometer (Tecan Systems Inc., San Jose, CA) set to 30°C with intermittent shaking, and  $\text{OD}_{595}$  measurements were taken at 15-min intervals for 24 hr. Raw absorbance data were averaged for all replicates, corrected for background, and plotted as a function of time. The area under the curve (AUC) was calculated for the cultures in each well using Prism software (version 5.01; GraphPad Software, Inc., La Jolla, CA), and the treatments were averaged and expressed as a percentage of the control.

**Human tissue array and real-time quantitative polymerase chain reaction (PCR) assay.** We used TaqMan-based real-time quantitative polymerase chain reaction (rt-qPCR) to quantify *N6AMT1* and *AS3MT* expression on a panel of 48 normal human tissues using the Human Rapid-Scan Plate (OriGene Technologies, Inc., Rockville, MD). The human tissues were selected from multiple individuals of different ethnicity and pooled together. We obtained the primers and probes used for amplification of *N6AMT1*, *AS3MT*, and *ACTB* ( $\beta$ -actin; control) from Applied Biosystems (Foster City, CA). Gene expression

of *N6AMT1* and *AS3MT* was calculated relative to *ACTB* using the  $\Delta\Delta C_T$  method.

***N6AMT1* gene expression vector constructs and stable cell lines.** Human *N6AMT1* cDNA (GenBank accession no. NM\_013240; National Center for Biotechnology Information 2011) was PCR amplified with primers 5'-AACGCAGCGAAGGACTAT-3' and 5'-CAGTAGTTCTGGGCACAC-3'. The PCR product was gel purified (Qiagen, Valencia, CA) and cloned into pcDNA 2.1 vector (Invitrogen) according to the manufacturer's instructions, and the sequence was confirmed. The pcDNA 2.1 vector containing the *N6AMT1* gene was excised using *NotI*/*Bam*HI restriction enzymes (New England Biolabs, Ipswich, MA) and subjected to gel purification. The nucleotides of the *N6AMT1* gene containing *Bam*HI and *NotI* overhangs were annealed and ligated to a linearized pRetro X-IRES-ZsGreen vector (Clontech, Mountain View, CA) digested with *Bam*HI and *NotI* (New England Biolabs). The pRetro X-IRES-ZsGreen vector is a fluorescent retroviral expression vector that allows both a gene of interest and the ZsGreen gene to be expressed. The resultant constructs were amplified, purified, and sequenced. UROtsa cells were transfected with this constructed vector or a control vector using Lipofectamine 2000 reagent (Invitrogen) according to the manufacturer's instructions. After incubation at 37°C for 8 hr, the supernatant fraction containing the retroviral vector was removed and replaced with normal growth medium. Cells grown for 48–72 hr were assessed by fluorescence microscopy. The ZsGreen fluorescent marker yields a bright green fluorescence, permitting direct monitoring of the delivery efficiency. Finally, the cell populations were sorted by the DAKO-Cytomation MoFlo High Speed Sorter (Dako North America, Carpinteria, CA), and the green fluorescent cells were purified and collected for continuing culture. The green fluorescent cells were used for additional experimentation.

**Semiquantitative reverse-transcription (RT)-PCR.** UROtsa cells with either *N6AMT1* or plasmid vectors were collected, and total RNA was isolated from these cells using the Qiagen RNeasy Mini kit. We performed a reverse transcription reaction using SuperScript II Reverse Transcriptase (Invitrogen) according to the manufacturer's instructions. The PCR conditions for DNA amplification in the linear range were established on the GeneAmp PCR System 7600 (PerkinElmer, Inc., Wellesley, MA). The primers for DNA amplification were as follows: *N6AMT1*, 5'-AACGCAGCGAAGGACTAT-3' and 5'-CAGTAGTTCTGGGCACAC-3'; *AS3MT*, 5'-GTGTCTGGGTGGTGCTT TATACTG-3' and 5'-TGGAGGGCAGA

ACCCAATT-3'; and the housekeeping gene *ACTB*, 5-TCACCCACACTGTGCCATCTACGA-3 and 5-CAGCGGAACCGCTCATTGCCAATGG-3. RT-PCR products were analyzed on 1% agarose gels.

**Cytotoxicity assay.** We performed the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) assay to assess cell viability after arsenic treatment. Cells were cultured in 96-well plates in a volume of 100 µL medium/well at a density of  $5 \times 10^4$  cells/mL. Twenty-four hours after incubation with  $\text{iAs}^{\text{III}}$  or  $\text{MMA}^{\text{III}}$  (six replicates/arsenic concentration), 10 µL sterile MTT dye (Sigma-Aldrich; 5 mg/mL) was added to each well and plates were incubated at 37°C for 4 hr. The culture medium was then removed, and 200 µL dimethyl sulfoxide was added and thoroughly mixed for 10 min. Spectrophotometric absorbance at 570 nm was measured in a microplate reader.

**Arsenic species profile analysis by high-performance liquid chromatography/inductively coupled plasma-mass spectrometry (HPLC-ICP-MS) methods.** UROtsa cells with *N6AMT1* and UROtsa cells with vector were grown in RPMI 1640 medium supplemented with 10% fetal bovine serum and antibiotics. Culture medium was collected after exposure to  $\text{iAs}^{\text{III}}$  or  $\text{MMA}^{\text{III}}$  for 24 hr or 3 days and stored at -80°C until analysis. Cells treated with  $\text{iAs}^{\text{III}}$  or  $\text{MMA}^{\text{III}}$  for 3 days were collected, lysed in RIPA buffer, and extracted in methanol by incubating overnight at 4°C in a rotational shaker. After centrifugation at maximum speed for 5 min at 4°C, supernatants were transferred to microcentrifuge tubes and stored at -80°C until analysis. Before analysis, samples were diluted 1:5 with water-methanol to bring the methanol concentration to 2.5%, incubated at 5°C to precipitate poorly soluble material, and filtered (0.45 µm). Analysis was performed by HPLC-ICP-MS (Agilent 1090 HPLC and Agilent 7500CE ICP-MS run in normal mode; both from Agilent Technologies, Santa Clara, CA) under conditions that resolved neutral, trivalent, and pentavalent  $\text{iAs}$  species. The ion-pairing method (Le et al. 2000) was used with major modifications to improve the resolution of the species. Briefly, calibrants were prepared from neat materials [ $\text{As}_2\text{O}_3$  (Aldrich) and  $\text{As}_2\text{O}_5$  (Acros), Sigma-Aldrich;  $\text{DMA}^{\text{V}}$  and  $\text{MMA}^{\text{V}} \cdot 6\text{H}_2\text{O}$ , Chem Service, West Chester, PA] in deionized water ( $\geq 18 \text{ M}\Omega$ ). For speciation, we used a Phenomenex Gemini-NX column (3 µm, C18, 110Å, 150 × 4.6 mm; Torrance, CA) with a corresponding guard column at 40°C. Concentrations of arsenic species in stock solutions were standardized against NIST traceable commercial ICP-MS standards (VWR BDH Aristar Plus; Ultra Scientific, Kingstown, RI). Serial dilutions were made into deionized water.  $\text{iAs}^{\text{III}}$  and  $\text{iAs}^{\text{V}}$  species were quantified by separate calibrant series, and  $\text{iAs}^{\text{III}}$  concentration in the

calibrants was corrected for any conversion to iAs<sup>V</sup>. The HPLC conditions were isocratic (5 mM tetrabutylammonium hydroxide, 10 mM ammonium carbonate, 2.5% methanol, pH 9.2, 1 mL/min) for 5 min; then a step gradient (5 mM tetrabutylammonium hydroxide, 30 mM ammonium carbonate, 2.5% methanol, pH 8.75, 1.2 mL/min) for 5 min to elute iAs<sup>V</sup> was followed by step gradients (5 mM tetrabutylammonium hydroxide, 30 mM ammonium carbonate, 2.5% methanol, pH 9.2, 1.2 mL/min) for a 5-min equilibration to the initial pH and finally to the initial mobile phase for 5 min (1.2 mL/min). The data were analyzed by LC ChemStation A.09.03 and ICP-MS ChemStation B.03.03 software (Agilent Technologies).

**Data analysis.** Statistical analyses were performed using one-way analysis of variance. Data represent mean  $\pm$  SE of at least three independent experiments.

## Results


**Deletion of yeast *MTQ2* leads to increased resistance to arsenic treatment.** We evaluated the growth phenotype of the *MTQ2*-deletion mutants in the presence of either iAs<sup>III</sup> or MMA<sup>III</sup> (Figure 1). Deletion strains and their isogenic wild-type counterpart, BY4743, were treated with equioxic doses equivalent to the concentrations that resulted in 20% growth inhibition (IC<sub>20</sub>) and 2  $\times$  IC<sub>20</sub>, which were 300 and 600  $\mu$ M for iAs<sup>III</sup> and 150 and

300  $\mu$ M for MMA<sup>III</sup>, respectively. iAs<sup>III</sup> treatments had no effect on the growth of *MTQ2*-deletion mutants but significantly decreased growth of the wild-type strain. In comparison, the growth of both *MTQ2*-deletion mutants and wild-type yeast treated with MMA<sup>III</sup> decreased to the same degree despite the higher toxicity of MMA<sup>III</sup>.

**Differential level of N6AMT1 mRNA expression in human tissues.** A direct search for sequence homology and conserved functional domains revealed that the human *N6AMT1* gene is orthologous to the yeast *MTQ2* gene. We used the web-based online tool Protein Function Prediction (PFP; Kihara Bioinformatics Laboratory 2010), to analyze and predict its potential functions (Hawkins et al. 2009). The suggested molecular functions of N6AMT1 include protein heterodimerization and methionine *S*-methyltransferase activity, with an almost certain (100%) predicted probability of being a methyltransferase. Given N6AMT1's suggested function as a methyltransferase, we were interested in exploring its potential involvement in arsenic biomethylation. Considering that the primary methyltransferase responsible for arsenic metabolism in human cells is AS3MT, we did pairwise alignment analysis of N6AMT1 and AS3MT using EMBOSS Pairwise Alignment Algorithms, an online tool (European Bioinformatics Institute 2010). The two

proteins shared about 25% similarity. Of the three sequence motifs found in most AS3MT homologs, only motif ILDLGSGSG is highly conserved in N6AMT1 [LEVGSGSG; see Supplemental Material, Figure 1 (doi:10.1289/ehp.1002733)], whereas (D/N)PPY is present in N6AMT1 but not in AS3MT. These differences suggest that the mechanism by which N6AMT1 methylates arsenic may differ from that of AS3MT, if N6AMT1 is, in fact, involved in the methylation of arsenicals.

A search of the Expressed Sequence Tags (EST) Database (National Center for Biotechnology Information 2010) revealed sequences matching the cDNA of *N6AMT1* in many human tissues with varied expression levels. To experimentally measure the mRNA expression of *N6AMT1* across tissues, we performed rt-qPCR analysis using cDNA from a panel of 48 human tissues contained in a tissue array (Figure 2). We confirmed the amplification products to be *N6AMT1* by DNA sequencing. Using the liver as a reference, the expression of *N6AMT1* was normalized to the expression level of *ACTB* [with cycle threshold (Ct) values ranging from 18 to 20] and found to be relatively highly expressed in tissues such as the parathyroid, pituitary, adrenal gland, and kidney, and weakly expressed in tissues such as the skin, lung, and mammary gland. We also measured *AS3MT* mRNA levels using the same tissue panel in order to compare *N6AMT1*




**Figure 1.** Deletion of the *MTQ2* gene in yeast results in increased resistance to arsenite (iAs<sup>III</sup>), shown as the growth phenotype of *MTQ2* mutant yeast cells (B) and the wild-type BY4743 cells (A) treated with 300 or 600  $\mu$ M iAs<sup>III</sup> or 150 or 300  $\mu$ M MMA<sup>III</sup>. Growth curves show the OD<sub>595</sub> for each treatment as a function of time for 24 hr. Bars represent the mean  $\pm$  SE AUC for three technical replicates. At the doses tested, iAs<sup>III</sup> treatment did not alter the growth pattern of *MTQ2* mutants but led to a dose-dependent reduction in growth of the wild-type strain; the growth patterns of both yeast strains were similar after MMA<sup>III</sup> exposure. #*p* < 0.001, compared with control.




expression in each tissue and *AS3MT* expression [see Supplemental Material, Figure 2 (doi:10.1289/ehp.1002733)]. The data showed that the level of *N6AMT1* mRNA

was relatively low in most measured tissues compared with *AS3MT* mRNA. The presence of detectable levels of *N6AMT1* expression in all tissues analyzed suggests that this enzyme

could be involved in methylating arsenicals. *N6AMT1* may act in parallel to *AS3MT* and may be functional only under certain conditions or in certain tissues.



**Figure 2.** *N6AMT1* mRNA expression in human tissues, quantified by rt-qPCR analysis a panel of 48 normal human tissues; transcript levels of *N6AMT1* were normalized to *ACTB* expression and are shown as the fold change relative to liver.



**Figure 3.** Enhancing expression of human *N6AMT1* in UROtsa cells increases resistance to both *iAs*<sup>III</sup> and *MMA*<sup>III</sup> treatment. (A) Representative images of UROtsa cells with either plasmid alone or plasmid with *N6AMT1*; bars = 10  $\mu$ m. (B) Semiquantitative RT-PCR analysis shows dramatically increased *N6AMT1* expression in UROtsa cells with *N6AMT1* relative to control cells or cells with plasmid. *AS3MT* mRNA was not detected in either cell line; *ACTB* was used as the loading control. UROtsa cells with vector and with *N6AMT1* were treated with increasing concentrations of *iAs*<sup>III</sup> (C) or *MMA*<sup>III</sup> (D) for 24 hr, and cell viability was evaluated with MTT. Bars represent the mean  $\pm$  SD of three independent experiments. Treatment with either *iAs*<sup>III</sup> or *MMA*<sup>III</sup> resulted in a dose-dependent decrease in viability. However, increased expression of *N6AMT1* in UROtsa cells led to resistance to both *iAs*<sup>III</sup> (C) and *MMA*<sup>III</sup> (D); this effect was more significant when cells were treated with *MMA*<sup>III</sup> than with *iAs*<sup>III</sup>.

\*\* $p < 0.01$ , compared with control and vector control.

**Overexpression of N6AMT1 in UROtsa cells increases resistance to arsenic treatment.** We also measured and compared the level of *N6AMT1* mRNA in several cell lines, including 293 (human embryonic kidney cells), HeLa, UROtsa, and HL60 (human promyelocytic leukemia cells). *N6AMT1* expression in UROtsa cells is relatively low, with a Ct value of about 33 (data not shown). This cell line also has almost no detectable level of *AS3MT*, making it an excellent model to study the role of *N6AMT1* in arsenic toxicity and metabolism in mammals. We enhanced *N6AMT1* expression in UROtsa cells using a retrovirus-based vector (Figure 3A) and found the level of *N6AMT1* mRNA in UROtsa cells to be significantly increased by approximately 5-fold in clone 2, as measured by semiquantitative RT-PCR (Figure 3B). We also measured *AS3MT* gene expression in these two cell lines and found no detectable mRNA level in either cell line (Figure 3B). We further confirmed these PCR results by real-time PCR analysis (data not shown). Unfortunately, we could not detect N6AMT1 protein levels in these cells using two commercially available antibodies. Transfected UROtsa cells did not have an altered doubling time or morphology in culture. The UROtsa cells with *N6AMT1* (*N6AMT1*-enhanced cells) and the UROtsa cells with vector cells (vector control) were treated with either iAs<sup>III</sup> or MMA<sup>III</sup> at a series of concentrations for 24 hr. Arsenical treatments induced a dose-dependent decrease in viability of both cell lines. However, increased expression of *N6AMT1* in UROtsa cells resulted in higher viability after iAs<sup>III</sup> and MMA<sup>III</sup> treatment at almost all concentrations tested, compared with the UROtsa vector control cells (Figure 3C,D). This increased arsenic resistance was more apparent in cultures treated with MMA<sup>III</sup> than with iAs<sup>III</sup>, approximately 2- and 1.3-fold, respectively.

**Enhanced N6AMT1 in UROtsa cells methylates MMA<sup>III</sup> to DMA.** We collected medium and cell extracts from cultures of UROtsa cells with *N6AMT1* and UROtsa cells with vector treated with different concentrations of either iAs<sup>III</sup> or MMA<sup>III</sup> for up to 3 days and then measured and analyzed arsenic metabolic profiles using ICP-MS [for representative chromatograms, see Supplemental Material, Figure 3 (doi:10.1289/ehp.1002733)]. Methylated metabolites were undetectable either in the media (Table 1) or in the cell extract (Table 2) from cultures of UROtsa cells with *N6AMT1* or of UROtsa cells with vector, after treatment of with iAs<sup>III</sup>, suggesting that N6AMT1 does not methylate iAs<sup>III</sup>. When UROtsa cells with vector control were treated with MMA<sup>III</sup>, levels of MMA<sup>V</sup> but not dimethylarsinic acid (DMA<sup>V</sup>) increased in the media (Table 3) and cells (Table 2). Treatment with MMA<sup>III</sup> of UROtsa cells with

*N6AMT1* resulted in similarly increased levels of MMA<sup>V</sup> and led to increased levels of DMA<sup>V</sup> in the media after 24 and 72 hr (Table 3) and, to a lesser degree, in the cell extracts after 72 hr (Table 2). At 24 hr, the level of DMA<sup>V</sup> was similar at each dose level, but after 3 days treatment levels in both media and cell extract increased in relation to the initial concentration of MMA<sup>III</sup>, in a dose-dependent manner. Moreover, the amount of DMA<sup>V</sup> in the culture medium increased 5-fold after 3 days of treatment (1 μM MMA<sup>III</sup>) compared with 1 day of treatment.

## Discussion

**MMA<sup>III</sup> is the most toxic arsenic metabolite in vivo and in vitro.** In humans, iAs is metabolized to methylated arsenical species in a multistep process. Methylated arsenicals,

especially MMA<sup>III</sup>, may be more toxic than iAs both *in vivo* and *in vitro* (Drobná et al. 2005; Ferrario et al. 2008; Kligerman et al. 2003; Petrick et al. 2001). In cultured human cells, MMA<sup>III</sup> is the most toxic arsenical (Drobná et al. 2005; Ferrario et al. 2008; Petrick et al. 2001) and inhibits several key cellular proteins, such as glutathione reductase (Styblo et al. 1997) and thioredoxin reductase (Lin et al. 1999, 2001). Several studies have shown that MMA<sup>III</sup> is capable of inducing genetic damage and changes in signal transduction by either direct or indirect mechanisms (Ahmad et al. 2002; Kligerman et al. 2003; Nesnow et al. 2002). In addition, exposure to MMA<sup>III</sup> for 52 weeks induced malignant transformation of UROtsa cells (Bredfeldt et al. 2006). Epidemiological studies have suggested that individuals who

**Table 1.** Arsenic metabolic profile in cell culture medium after iAs<sup>III</sup> treatment of UROtsa cells with vector and UROtsa cells with *N6AMT1* (ng/mL; mean ± SE).

iAs <sup>III</sup> (μM)	UROtsa cells with vector				UROtsa cells with <i>N6AMT1</i>			
	iAs <sup>III(+V)</sup>	MMA <sup>III</sup>	MMA <sup>V</sup>	DMA <sup>V</sup>	iAs <sup>III(+V)</sup>	MMA <sup>III</sup>	MMA <sup>V</sup>	DMA <sup>V</sup>
<b>1 day</b>								
0	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ
1	261.50 ± 126.22	BLOQ	BLOQ	BLOQ	417.00 ± 1.41	BLOQ	BLOQ	BLOQ
3	1109.00 ± 11.31	BLOQ	BLOQ	BLOQ	1159.00 ± 12.02	BLOQ	BLOQ	BLOQ
10	3695.50 ± 10.96	BLOQ	BLOQ	BLOQ	3751.00 ± 72.12	BLOQ	BLOQ	BLOQ
<b>3 days</b>								
0	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ
1	425.50 ± 8.13	BLOQ	BLOQ	BLOQ	414.50 ± 5.30	BLOQ	BLOQ	BLOQ
3	1034.00 ± 2.12	BLOQ	BLOQ	BLOQ	1017.00 ± 7.07	BLOQ	BLOQ	BLOQ

BLOQ, below the limit of quantitation (the *m/z* 75 signal at the retention time of the species is not distinguishable from the baseline signal).

**Table 2.** Arsenic metabolic profile in cell extract after 3-day iAs<sup>III</sup> and MMA<sup>III</sup> treatment of UROtsa cells with vector and UROtsa cells with *N6AMT1* (ng/mL; mean ± SE).

Treatment (μM)	UROtsa cells with vector				UROtsa cells with <i>N6AMT1</i>			
	iAs <sup>III(+V)</sup>	MMA <sup>III</sup>	MMA <sup>V</sup>	DMA <sup>V</sup>	iAs <sup>III(+V)</sup>	MMA <sup>III</sup>	MMA <sup>V</sup>	DMA <sup>V</sup>
<b>iAs<sup>III</sup></b>								
0	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ
1	2.7 ± 0.3	BLOQ	BLOQ	BLOQ	5.7 ± 0.6	BLOQ	BLOQ	BLOQ
3	12.0 ± 2.0	BLOQ	BLOQ	BLOQ	14.5 ± 0.5	BLOQ	BLOQ	BLOQ
<b>MMA<sup>III</sup></b>								
0	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ
0.5	BLOQ	BLOQ	3.5 ± 0.7	BLOQ	BLOQ	BLOQ	3.0 ± 0.1	2.8 ± 0.2
1	BLOQ	BLOQ	2.9 ± 0.8	BLOQ	BLOQ	BLOQ	4.7 ± 0.0	4.8 ± 0.6

BLOQ, below the limit of quantitation (the *m/z* 75 signal at the retention time of the species is not distinguishable from the baseline signal).

**Table 3.** Arsenic metabolic profile in cell culture medium after MMA<sup>III</sup> treatment of UROtsa cells with vector and UROtsa cells with *N6AMT1* (ng/mL; mean ± SE).

MMA <sup>III</sup> (μM)	UROtsa cells with vector				UROtsa cells with <i>N6AMT1</i>			
	iAs <sup>III(+V)</sup>	MMA <sup>III</sup>	MMA <sup>V</sup>	DMA <sup>V</sup>	iAs <sup>III(+V)</sup>	MMA <sup>III</sup>	MMA <sup>V</sup>	DMA <sup>V</sup>
<b>Treated 1 day</b>								
0	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ
0.5	BLOQ	Present	112.00 ± 0.71	BLOQ	BLOQ	Present	104.00 ± 0.71	21.50 ± 0.35
1	BLOQ	Present	236.50 ± 15.20	BLOQ	BLOQ	Present	210.00 ± 2.12	22.50 ± 1.06
2	BLOQ	Present	468.00 ± 17.68	BLOQ	BLOQ	Present	424.50 ± 6.72	20.00 ± 0.71
<b>Treated 3 days</b>								
0	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ
0.5	BLOQ	Present	115.00 ± 2.12	BLOQ	BLOQ	Present	106.00 ± 2.83	72.00 ± 4.24
1	BLOQ	Present	277.00 ± 5.66	BLOQ	BLOQ	Present	236.50 ± 9.55	105.50 ± 4.60

BLOQ, below the limit of quantitation (the *m/z* 75 signal at the retention time of the species is not distinguishable from the baseline signal). "Present" indicates that the peak was seen at MMA<sup>III</sup> retention time; no concentration is given because the samples were not analyzed by the MMA<sup>III</sup> assay.

excrete a higher proportion of ingested arsenic as MMA<sup>III</sup> are more susceptible to arsenic-related cancer (Steinmaus et al. 2006). MMA<sup>III</sup> has been proposed as the ultimate genotoxic form of arsenic (Kligerman et al. 2003), and the existing evidence indicates that biomethylation of iAs to MMA<sup>III</sup> is likely to alter the adverse effects of environmental arsenic exposure on human health.

*AS3MT is primarily responsible for methylating iAs to MMA<sup>III</sup> and DMA in humans.* The arsenic methyltransferase AS3MT is recognized as the primary enzyme responsible for conversion of iAs to its methylated metabolites MMA<sup>III</sup> and DMA (Lin et al. 2002; Wood et al. 2006). Studies have shown that single-nucleotide polymorphisms in *AS3MT* lead to different urinary arsenical profiles (Agusa et al. 2009; Schläwicke Engström et al. 2009; Wood et al. 2006), some of which are associated with increased risk of premalignant skin lesions (Valenzuela et al. 2009). Although these data suggest that AS3MT plays a critical role in arsenic methylation and toxicity, a recent study showed that *As3mt*-knockout mice retain some ability to methylate arsenicals, suggesting the existence of other methyltransferases that could be involved in alternative arsenic metabolism pathways (Drobna et al. 2009).

*N6AMT1 is capable of methylating MMA<sup>III</sup> to DMA.* In this study, we found that N6AMT1 has the capacity to methylate MMA<sup>III</sup> to DMA. The expression of *N6AMT1* is generally low compared with the expression level of *AS3MT* in most human tissues. In addition, the low sequence homology shared between these two proteins, about 25%, supports differences in substrate specificity and, possibly, in mechanisms of arsenic methylation. In contrast to AS3MT, which methylates iAs to the more toxic MMA<sup>III</sup>, N6AMT1 methylates MMA<sup>III</sup> to the less toxic DMA. This is consistent with the increased resistance to MMA<sup>III</sup> of UROtsa cells overexpressing *N6AMT1* compared with vector control cells, but other mechanisms likely also contribute to this increased resistance. Thus, our results suggest that *N6AMT1* may play a role in modulating arsenical-induced toxicity and that decreased *N6AMT1* expression or activity could have a significant impact on arsenic-induced toxicity and perhaps carcinogenicity under certain conditions or in certain tissues.

*N6AMT1 in human cells responded to arsenicals differently from MTQ2 in yeast.* We noted differences in response to arsenical treatments between yeast and human cells. Specifically, the *MTQ2*-deletion yeast strain is resistant only to iAs<sup>III</sup>. We did not find evidence of iAs<sup>III</sup> methylation in yeast wild-type; that is, DMA and MMA levels in cells and culture media were below the limit of quantitation (data not shown). Therefore, the function of

*MTQ2* in arsenic toxicity may not be related to iAs<sup>III</sup> methylation. In contrast, overexpression of *N6AMT1* in UROtsa cells leads to resistance to both iAs<sup>III</sup> and MMA<sup>III</sup>, which provides evidence of its involvement in protection from these arsenicals. These results indicate that the orthologous genes—*MTQ2* in yeast and *N6AMT1* in humans—have different roles in the cellular response to arsenic toxicity.

*Conversion of MMA<sup>III</sup> to DMA by N6AMT1 needs to be further confirmed biochemically.* Our analyses showed that enhanced expression of *N6AMT1* in UROtsa cells converts MMA to DMA. However, purified recombinant N6AMT1, in the presence of SAM and other cofactors, was unable to methylate iAs<sup>III</sup> or MMA<sup>III</sup> (data not shown). N6AMT1 dimerizes with tRNA methyltransferase 11-2 homolog (TRMT112), which appears to be necessary for proper N6AMT1 function (Figaro et al. 2008) and is consistent with heterodimerization activity as predicted with PFP. Thus, the lack of N6AMT1-dependent MMA<sup>III</sup> methylation in this test tube experiment may be due to the absence of TRMT112 or another unknown protein. In addition, *N6AMT1* overexpression increased resistance to iAs<sup>III</sup>, although UROtsa cells were not able to methylate iAs<sup>III</sup> to MMA<sup>III</sup> or to other methylated species, suggesting that mechanisms other than participation in arsenic methylation might be involved. It is not clear whether interaction between N6AMT1 and TRMT112 has a role in arsenic toxicity, but it is certainly worthy of further investigation.

## Conclusions

Our data suggest an important potential role of *N6AMT1* in modulating arsenical-induced toxicity by methylating MMA<sup>III</sup> to the less toxic DMA. However, further investigation is warranted to determine whether N6AMT1 can methylate MMA<sup>III</sup> *in vivo* and also to identify the genetic and environmental factors that can alter *N6AMT1* expression and/or activity. Our ongoing experiments are focused on the biochemical characterization of N6AMT1, specifically its capacity to methylate MMA<sup>III</sup>, as well as its ability to modulate arsenic toxicity and carcinogenicity.

## REFERENCES

- Abernathy CO, Liu YP, Longfellow D, Aposhian HV, Beck B, Fowler B, et al. 1999. Arsenic: health effects, mechanisms of actions, and research issues. *Environ Health Perspect* 107:593–597.
- Agusa T, Iwata H, Fujihara J, Kunito T, Takeshita H, Minh TB, et al. 2009. Genetic polymorphisms in AS3MT and arsenic metabolism in residents of the Red River Delta, Vietnam. *Toxicol Appl Pharmacol* 236(2):131–141.
- Ahmad S, Kitchin KT, Cullen WR. 2002. Plasmid DNA damage caused by methylated arsenicals, ascorbic acid and human liver ferritin. *Toxicol Lett* 133(1):47–57.
- Bentley R, Chasteen TG. 2002. Microbial methylation of metalloids: arsenic, antimony, and bismuth. *Microbiol Mol Biol Rev* 66(2):250–271.
- Bredfeldt TG, Jagadish B, Eblin KE, Mash EA, Gandolfi AJ. 2006. Monomethylarsonous acid induces transformation of human bladder cells. *Toxicol Appl Pharmacol* 216(1):69–79.
- Drobna Z, Naranmandura H, Kubachka KM, Edwards BC, Herbin-Davis K, Styblo M, et al. 2009. Disruption of the arsenic (+3 oxidation state) methyltransferase gene in the mouse alters the phenotype for methylation of arsenic and affects distribution and retention of orally administered arsenate. *Chem Res Toxicol* 22(10):1713–1720.
- Drobna Z, Waters SB, Devesa V, Harmon AW, Thomas DJ, Styblo M. 2005. Metabolism and toxicity of arsenic in human urothelial cells expressing rat arsenic (+3 oxidation state)-methyltransferase. *Toxicol Appl Pharmacol* 207(2):147–159.
- Eblin KE, Bredfeldt TG, Gandolfi AJ. 2008. Immortalized human urothelial cells as a model of arsenic-induced bladder cancer. *Toxicology* 248(2–3):67–76.
- European Bioinformatics Institute. 2010. EMBOS Pairwise Alignment Algorithms. Available: <http://www.ebi.ac.uk/Tools/emboss/align/index.html> [accessed 15 October 2010].
- Ferrario D, Croera C, Brustio R, Collotta A, Bowe G, Vahter M, et al. 2008. Toxicity of inorganic arsenic and its metabolites on haematopoietic progenitors "in vitro": comparison between species and sexes. *Toxicology* 249(2–3):102–108.
- Figaro S, Scrima N, Buckingham RH, Heurgue-Hamard V. 2008. HemK2 protein, encoded on human chromosome 21, methylates translation termination factor eRF1. *FEBS Lett* 582(16):2352–2356.
- Hall LL, George SE, Kohan MJ, Styblo M, Thomas DJ. 1997. In vitro methylation of inorganic arsenic in mouse intestinal cecum. *Toxicol Appl Pharmacol* 147(1):101–109.
- Hawkins T, Chitale M, Luban S, Kihara D. 2009. PFP: Automated prediction of gene ontology functional annotations with confidence scores using protein sequence data. *Proteins* 74(3):566–582.
- IARC (International Agency for Research on Cancer). 2004. Some Drinking-water Disinfectants and Contaminants, including Arsenic. IARC Monogr Eval Carcinog Risks Hum 84:1–477.
- Jo WJ, Loguinov A, Wintz H, Chang M, Smith AH, Kalman D, et al. 2009. Comparative functional genomic analysis identifies distinct and overlapping sets of genes required for resistance to monomethylarsonous acid (MMA<sup>III</sup>) and arsenite (As<sup>III</sup>) in yeast. *Toxicol Sci* 111(2):424–436.
- Kapaj S, Peterson H, Liber K, Bhattacharya P. 2006. Human health effects from chronic arsenic poisoning—a review. *J Environ Sci Health B* 41(10):2399–2428.
- Kihara Bioinformatics Laboratory. 2010. PFP: Automated Protein Function Prediction Job Submission. Available: <http://dragon.bio.purdue.edu/pfp/> [accessed 15 October 2010].
- Kligerman AD, Doerr CL, Tennant AH, Harrington-Brock K, Allen JW, Winkfield E, et al. 2003. Methylated trivalent arsenicals as candidate ultimate genotoxic forms of arsenic: induction of chromosomal mutations but not gene mutations. *Environ Mol Mutagen* 42(3):192–205.
- Le XC, Ma M, Cullen WR, Aposhian HV, Lu X, Zheng B. 2000. Determination of monomethylarsonous acid, a key arsenic methylation intermediate, in human urine. *Environ Health Perspect* 108:1015–1018.
- Lin S, Cullen WR, Thomas DJ. 1999. Methylarsenicals and arsinothiolates are potent inhibitors of mouse liver thioredoxin reductase. *Chem Res Toxicol* 12(10):924–930.
- Lin S, Del Razo LM, Styblo M, Wang C, Cullen WR, Thomas DJ. 2001. Arsenicals inhibit thioredoxin reductase in cultured rat hepatocytes. *Chem Res Toxicol* 14(3):305–311.
- Lin S, Shi Q, Nix FB, Styblo M, Beck MA, Herbin-Davis KM, et al. 2002. A novel S-adenosyl-L-methionine:arsenic(III) methyltransferase from rat liver cytosol. *J Biol Chem* 277(13):10795–10803.
- National Center for Biotechnology Information. 2010. Expressed Sequence Tags Database. Available: <http://www.ncbi.nlm.nih.gov/dbEST/> [accessed 15 October 2010].
- National Center for Biotechnology Information. 2011. GenBank. Available: <http://www.ncbi.nlm.nih.gov/sites/entrez?db=nucleotide> [accessed 15 October 2010].
- Nesnow S, Rupp BC, Lambert G, Kadiiska M, Mason RP, Cullen WR, et al. 2002. DNA damage induced by methylated trivalent arsenicals is mediated by reactive oxygen species. *Chem Res Toxicol* 15(12):1627–1634.
- Pershagen G. 1981. The carcinogenicity of arsenic. *Environ Health Perspect* 40:93–100.
- Petrick JS, Jagadish B, Mash EA, Aposhian HV. 2001. Monomethylarsonous acid (MMA<sup>III</sup>) and arsenite: LD<sub>50</sub> in hamsters and in vitro inhibition of pyruvate dehydrogenase. *Chem Res Toxicol* 14(6):651–656.
- Polevoda B, Span L, Sherman F. 2006. The yeast translation

- release factors Mrf1p and Sup45p (eRF1) are methylated, respectively, by the methyltransferases Mttq1p and Mttq2p. *J Biol Chem* 281(5):2562–2571.
- Ratel D, Ravanat JL, Charles MP, Pladet N, Breuillaud L, Lunardi J, et al. 2006. Undetectable levels of N6-methyl adenine in mouse DNA: cloning and analysis of PRED28, a gene coding for a putative mammalian DNA adenine methyltransferase. *FEBS Lett* 580(13):3179–3184.
- Schläwicke Engström K, Nermell B, Concha G, Strömberg U, Vahter M, Broberg K. 2009. Arsenic metabolism is influenced by polymorphisms in genes involved in one-carbon metabolism and reduction reactions. *Mutat Res* 667(1–2):4–14.
- Sens DA, Park S, Gurel V, Sens MA, Garrett SH, Somji S. 2004. Inorganic cadmium- and arsenite-induced malignant transformation of human bladder urothelial cells. *Toxicol Sci* 79(1):56–63.
- Smith AH, Hopenhayn-Rich C, Bates MN, Goeden HM, Hertz-Picciotto I, Duggan HM, et al. 1992. Cancer risks from arsenic in drinking water. *Environ Health Perspect* 97:259–267.
- Smith AH, Steinmaus CM. 2009. Health effects of arsenic and chromium in drinking water: recent human findings. *Annu Rev Public Health* 30:107–122.
- Steinmaus C, Bates MN, Yuan Y, Kalman D, Atallah R, Rey OA, et al. 2006. Arsenic methylation and bladder cancer risk in case-control studies in Argentina and the United States. *J Occup Environ Med* 48(5):478–488.
- Stephens C, Reisenauer A, Wright R, Shapiro L. 1996. A cell cycle-regulated bacterial DNA methyltransferase is essential for viability. *Proc Natl Acad Sci USA* 93(3):1210–1214.
- Straif K, Benbrahim-Tallaa L, Baan R, Grosse Y, Secretan B, El Ghissassi F, et al. 2009. A review of human carcinogens—part C: metals, arsenic, dusts, and fibres. *Lancet Oncol* 10:453–454.
- Stybło M, Del Razo LM, Vega L, Germolec DR, LeCluyse EL, Hamilton GA, et al. 2000. Comparative toxicity of trivalent and pentavalent inorganic and methylated arsenicals in rat and human cells. *Arch Toxicol* 74(6):289–299.
- Stybło M, Serves SV, Cullen WR, Thomas DJ. 1997. Comparative inhibition of yeast glutathione reductase by arsenicals and arsenothiols. *Chem Res Toxicol* 10(1):27–33.
- Thomas DJ, Li J, Waters SB, Xing W, Adair BM, Drobná Z, et al. 2007. Arsenic (+3 oxidation state) methyltransferase and the methylation of arsenicals. *Exp Biol Med (Maywood)* 232(1):3–13.
- Valenzuela OL, Drobná Z, Hernández-Castellanos E, Sánchez-Peña LC, García-Vargas GG, Borja-Aburto VH, et al. 2009. Association of AS3MT polymorphisms and the risk of pre-malignant arsenic skin lesions. *Toxicol Appl Pharmacol* 239(2):200–207.
- Wood TC, Salavagionne OE, Mukherjee B, Wang L, Klumpp AF, Thomae BA, et al. 2006. Human arsenic methyltransferase (AS3MT) pharmacogenetics: gene resequencing and functional genomics studies. *J Biol Chem* 281(11):7364–7373.