

Pathways for the Anaerobic Microbial Debromination of Polybrominated Diphenyl Ethers

KRISTIN R. ROBROCK,[†] PETER KORYTÁR,[‡]
AND LISA ALVAREZ-COHEN^{*,†,§}

Department of Civil and Environmental Engineering,
University of California, Berkeley, California, Wageningen
IMARES-Institute for Marine Resources and Ecosystem Studies,
IJmuiden, The Netherlands, and Earth Sciences Division,
Lawrence Berkeley National Laboratory, Berkeley, California

Received August 21, 2007. Revised manuscript received
January 14, 2008. Accepted January 23, 2008.

The debromination pathways of seven polybrominated diphenyl ethers (PBDEs) by three different cultures of anaerobic dehalogenating bacteria were investigated using comprehensive two-dimensional gas chromatography (GC × GC). The congeners analyzed were the five major components of the industrially used octa-BDE mixture (octa-BDEs 196, 203, and 197, hepta-BDE 183, and hexa-BDE 153) as well as the two most commonly detected PBDEs in the environment, penta-BDE 99 and tetra-BDE 47. Among the dehalogenating cultures evaluated in this study were a trichloroethene-enriched consortium containing multiple *Dehalococcoides* species, and two pure cultures, *Dehalobacter restrictus* PER-K23 and *Desulfotobacterium hafniense* PCP-1. PBDE samples were analyzed by GC × GC coupled to an electron capture detector to maximize separation and identification of the product congeners. All studied congeners were debrominated to some extent by the three cultures and all exhibited similar debromination pathways with preferential removal of para and meta bromines. Debromination of the highly brominated congeners was extremely slow, with usually less than 10% of nM concentrations of PBDEs transformed after three months. In contrast, debromination of the lesser brominated congeners, such as penta 99 and tetra 47, was faster, with some cultures completely debrominating nM levels of tetra 47 within weeks.

Introduction

Polybrominated diphenyl ethers (PBDEs) are flame retardants that have been integrated into plastics and textiles for about three decades (1). Recent studies found that the penta-substituted BDEs are endocrine disruptors, thus raising concern over their elevated levels of detection in the environment and particularly in human breast milk (2, 3). Concentrations in environmental samples and human tissues have reportedly been increasing exponentially over time, doubling approximately every five years (4). A recent study of sediments from the San Francisco Bay Area reported PBDE concentrations up to 200 ppb (5). The penta- and octa-

brominated BDEs have been banned in the European Union and in several states in the United States and their production has ceased. However, the fully brominated deca-BDE continues to be used (6).

PBDEs can be debrominated under anaerobic conditions by dehalogenating bacteria (7) (8). Less brominated congeners, such as penta- and tetra-BDEs, are more toxic and bioavailable than more highly brominated congeners, such as octa-BDEs (9). It is therefore important to understand the dominant debromination pathways in order to predict the likely congeners produced from the transformation of environmentally prevalent PBDEs. Although specific rates of debromination determined in the laboratory can not be directly extrapolated to the environment, nonetheless, improved understanding of preferential transformation pathways can provide important information about the relative biodegradability or recalcitrance of individual PBDE congeners.

We previously found that a commercial octa-BDE mixture could be reductively debrominated to a variety of less brominated congeners including penta-BDE 99 and tetra-BDE 47 (8). Because of the complexity of the numerous starting substrates present in the octa mixture and the diversity of products that were formed, it was not possible to delineate specific PBDE debromination pathways in that study. To clearly identify specific debromination pathways, products generated by bacteria exposed to individual PBDE congeners must be identified. There are analytical challenges associated with PBDE congener identification since many of them (up to 50%) coelute when analyzed by conventional gas chromatography. Comprehensive two-dimensional chromatography employs two different and sequential chromatography columns that dramatically reduces the number of coelutions, significantly improving the separation and identification of PBDE congeners (10).

Our goal was to determine the debromination pathway of seven environmentally relevant PBDE congeners by a variety of dehalogenating bacteria. The congeners studied were those that dominate the industrial octa-BDE mixture DE-79, specifically octa 196, octa 203, octa 197, hepta 183, and hexa 153. In addition, two of the most frequently detected congeners in the environment, penta 99 and tetra 47, were included in this study. Bacteria representing three different dehalogenating genera, *Dehalococcoides*, *Dehalobacter*, and *Desulfotobacterium*, were evaluated to provide a broad understanding of anaerobic PBDE debromination. Our results describe the debromination pathways and the PBDE products that might be expected to occur in the environment under anaerobic conditions. In addition, this work represents the first use of comprehensive two-dimensional gas chromatography to understand PBDE debromination processes.

Materials and Methods

Chemicals. Individual congeners purchased from Accustandard (New Haven, CT) or Wellington Laboratories (Guelph, ON, Canada) were dissolved in nonane at a concentration of 50 µg/L. The octa-BDE mixture was obtained from Sigma-Aldrich (Sigma-Aldrich, Inc., St. Louis, MO). The major congeners in the octa-BDE mixture are deca (209), two nona congeners (207 and 206), three octa congeners (196, 203, and 197), three hepta congeners (183, 180, and 171), one hexa congener (153), and additional trace congeners (hexas 154 and 144; heptas 140, 138, and 191; octa 201 and 194; nona 208) (11). Decabromobiphenyl (DBB) and isooctane were purchased from Fisher Scientific (Fairlawn, NJ).

* Corresponding author phone: 510-643-5969; fax: 510-642-7483; e-mail: alvarez@ce.berkeley.edu.

[†] University of California, Berkeley.

[‡] Wageningen IMARES.


[§] Lawrence Berkeley National Laboratory.

Cultures. Three cultures were investigated in this study: (1) a trichloroethene-dechlorinating enrichment culture containing multiple *Dehalococcoides* strains (12, 13) mixed in a 1:10 ratio with the tetrachloroethene (PCE) dechlorinating strain *D. ethenogenes* 195 to form a mixed culture called ANAS195 (strain 195 was kindly donated by Professor Zinder at Cornell University (14)); (2) a PCE dechlorinating bacterium *Dehalobacter restrictus* PER-K23 (DSM 9455) (15); and (3) a pentachlorophenol degrading bacterium *Desulfotobacterium hafniense* PCP-1 (DSM 12420) (16), both obtained from the German Collection of Microorganisms and Cell Cultures. Additionally, we tested *Desulfotobacterium chlororespirans* Co23 (DSM 11544) (17), *Desulfotobacterium dehalogenans* JW/IU-DC1 (DSM 9161) (18), and *Desulfomonile tiedjei* DCB-1 (DSM 6799) (19) for their ability to debrominate the octa-BDE mixture.

Culture Media and Cultivation. Cultures were grown in 160 mL serum bottles with 100 mL of medium and sealed with butyl rubber septa (Bellco Glass Inc., Vineland, NJ) and aluminum crimp caps (Wheaton Science Products, Millville, NJ) to ensure a robust anaerobic environment over the course of the experiment. ANAS195 was grown in mineral salts medium (20) reduced with 0.2 mM cysteine sulfide with 20 mM lactate and 685 μ M PCE. *D. restrictus* was grown in the same medium as ANAS195 with 5 mM acetate, 685 μ M PCE, and 100 mg/L of peptone. The headspace of the bottles was H_2/CO_2 (80:20 v/v). *D. hafniense*, *D. dehalogenans*, and *D. chlororespirans* were grown in DSMZ medium 720 reduced with 1.25 mM Na_2S with 5 μ M pentachlorophenol, 10 mM 3-chloro-4-hydroxyphenylacetate, and 10 mM 3-chloro-4-hydroxybenzoate, respectively. *D. tiedjei* was grown in DSMZ medium 521 with 10 mM 3-chlorobenzoic acid. Octa-BDE was added to these cultures dissolved in nonane for final concentrations of 1 μ M. Individual PBDE congeners dissolved in nonane were added to the samples for a final concentration of 20 μ g/L (25 nM for the octa-BDEs, 27 nM hepta-BDE 183, 30 nM hexa-BDE 153, 35 nM penta-BDE 99, and 40 nM tetra-BDE 47). All bottles contained resazurin as an oxygen indicator. Active or autoclaved cultures were inoculated (10% v/v for the ANAS195 culture and 5% v/v for all other cultures) after addition of the PBDEs and complete dissolution of the chlorinated solvents. Uninoculated abiotic controls were used for the spore-forming *D. hafniense* as original autoclaved controls for this culture revived during incubation. All samples and controls were incubated at 30 °C in the dark without shaking. Experiments were conducted with triplicate biological samples and single controls and were monitored for three months. Most experiments were repeated for verification of the results.

Sample Preparation. Monthly, 1 mL of culture was removed from the samples by disposable plastic syringe (Becton, Dickinson and Company, Franklin Lakes, NJ). A 2- μ L portion of methanol saturated with decabromobiphenyl (DBB) was added to the samples as an internal standard to determine extraction efficiency. Surrogate extraction efficiencies ranged from 77 to 82%. The mixture was subjected to a liquid-liquid extraction with 1 mL of isooctane in a 4-mL amber glass vial. The mixture was vortexed for 2 min and shaken for 16 h. Samples were centrifuged at 15 800g for 3 min to remove particulate matter and bacterial debris. The solvent phase isooctane (\sim 950 μ L) was transferred to a 2-mL amber glass vial for analysis. Samples were concentrated 10-fold with a stream of nitrogen to improve the limit of detection.

Analytical Methods. PBDE congeners were detected using comprehensive two-dimensional gas chromatography with electron capture detection at the Wageningen IMARES laboratories in IJmuiden, Netherlands (10). Two-dimensional chromatography uses two different columns in series to separate compounds based on differing physical properties






FIGURE 2. Debromination of Hepta 183 by the ANAS195 culture. The GC \times GC chromatograms show (A) the autoclaved control and (B) an active sample showing generation of various hexa-BDE congeners during debromination of Hepta-BDE 183 after 3 months.

at 20 °C/min to 150 °C, increased at 2 °C/min until 325 °C where it was held for 15 min. Injector and detector temperatures were 250 and 350 °C respectively. A 2- μ L portion of the sample was injected into a split/splitless inlet port operated in the splitless mode with split opening 2 min after injection. The identification of specific PBDEs was performed by comparing two-dimensional peak retention times with standard solutions containing 138 mono through deca-BDE congener standards. PBDE concentrations were calculated using calibration curves specific for each congener between 0.5 and 30 nM. Average concentrations curves for homologue groups were used only for the congeners for which standards were not available. The instrument detection limits for PBDE congeners were 0.1–1.0 fmol per injection, corresponding to low pM concentrations in the samples. To account for different extraction efficiencies over the long time course of these experiments, the amount of PBDE debromination reported here is presented as the molar fraction of the total amount of recovered PBDEs.

Cell Counts. For enumeration of bacterial cells, real-time quantitative polymerase chain reaction (qPCR) was performed on the 16S rRNA gene as described previously (21). Cell DNA was extracted using a MoBio Microbial DNA Isolation kit. The following primer pair and probe specific to *Dehalococcoides* spp. were employed: forward primer 5'-GGTAATACGTAGGGAAGCAAGCG-3', probe 5'-VIC-ACATC-CAACTTGAAAGACCACCTACGCTCACT-TAMRA-3', reverse primer 5'-CCGGTTAAGCCGGGAAATT-3' (13). Primers designed for *D. restrictus* PER-K23 were 5'-CGG AGA TAG TTA ATG AAG CTT GCG -3' forward and 5'-TTT CCA GCT GTT GTC CCG GT -3' reverse and for *D. hafniense* PCP-1 were 5'-CGA ACG GTC CAG TGT CTA -3' forward and 5'-ACT TAT TGA ATG CGT GGA GCG -3' reverse (22). SYBR Green was used instead of specific probes to quantify both *D. restrictus* and *D. hafniense*. Calibration curves ranged from 10^3 to 10^8 cells/mL.

Results

We began by exposing a variety of anaerobic dehalogenating bacterial cultures to the industrial octabromodiphenyl ether mixture DE-79 to determine if they could debrominate PBDEs. An enrichment culture containing multiple *Dehalococcoides* species, designated ANAS195, produced a variety of debromination congeners when exposed to the octa-BDE mixture (8). Similar results were obtained with *Desulfotobacterium hafniense* PCP-1, *Desulfotobacterium chlororespirans* Co23, *Desulfotobacterium dehalogenans* JW/IU-DC1, and *Dehalobacter restrictus* PER-K23 (data not shown). No debromination of the octa-BDE mixture occurred with *Desulfomonile tiedjei* DCB-1. When the ANAS195 samples were analyzed with comprehensive two-dimensional gas chromatography they revealed numerous previously undetected dehalogenation products. Figure 1c shows the wide variety of debromination products produced from the octa-BDE mixture by the ANAS195 consortium. The products range from hepta to tribrominated congeners. With the improved separation enabled by GC \times GC, a number of peaks could be matched to PBDE standards and identified, including penta-BDE 99 and tetra-BDE 47—two of the most toxic PBDE congeners (23, 24).

The complexity of multiple starting substrates and products generated from transformation of the octa-BDE mixture precluded the identification of specific PBDE debromination pathways. Three diverse and representative debrominating cultures, ANAS195, *D. restrictus*, and *D. hafniense*, were therefore exposed individually to the five major components of the octa-BDE mixture (octas 196, 203, and 197; hepta 183, and hexa 153). Additionally, two environmentally important congeners that had been detected as products of the octa-BDE mixture transformation (penta 99 and tetra 47) were also tested. As a representation, dehalogenation products of hepta-BDE 183 by the ANAS195 culture are shown in Figure 2. Product peaks are clearly visible in the live sample that are not present in the autoclaved control.

Since the objective of this study was identification of debromination pathways rather than the calculation of biotransformation kinetics, and because the likely variability in extraction efficiencies for the wide variety of individual PBDE congeners detected in this study could not be adequately quantified using the single internal standard method employed herein, mass balances on debromination products were not calculated. Therefore, debromination products in Table 1 are represented as percent of the total molar concentration of recovered PBDEs. Some products were present only in trace quantities and are listed as detectable but not quantifiable (DNQ).

All tested congeners were found to be biotransformable by all cultures within three months of incubation, although

TABLE 1. Products Generated by the Debromination of Individual PBDE Congeners by Microbial Culture after Three Months^a

culture	octa 197	octa 203	octa 196	hepta 183	hexa 153	penta 99	tetra 47
ANAS195	hepta 184 0.8% (0.5–1.1%)	hepta 187* 2.6% (2.3–3.0%)	hepta 183 0.8% (0.4–1.5%)	hexa 154 4.1% (3.8–4.8%)	penta 118 0.6% (0.2–0.7%)	tetra 66 0.1% (0.1–0.2%)	tri 28 0.2% (0.17–0.2%)
	hepta 183 0.6% (0.4–0.9%)	hepta 183 2.2% (2.0–2.5%)	hepta 182 0.5% (0.2–1%)	hexa 153 2.3% (2.0–2.9%)	penta 101 0.6% (0.3–0.9%)	tetra 49 0.6% (0.4–1%)	tri 17 5.0% (4.3–5.6%)
	hepta 176 0.4% (0.3–0.7%)	hepta 180 0.3% (0.25–0.3%)	hepta 180 0.3% (0.1–0.6%)	hexa 149 3.3% (2.9–3.5%)	penta 99 0.9% (0.8–1.0%)	tetra 47 0.2% (0.2–0.3%)	
			Hepta 175* 1.2% (0.4–2.8%)	hexa 144 0.1% (0.0–0.3%)	tetra 47 (DNQ)		
				penta 100 (DNQ)			
<i>D. hafniense</i>	hepta 184 0.3% (0.1%–0.4%)	hepta 187* 2.3% (1.4–3.3%)	hepta 183 3.4% (1.7–5.1%)	hexa 154 7.1% (5.7–9.7%)	penta 101 10.3% (6.8–21%)	tetra 66 (DNQ)	tri 17 44% (17–71%)
	hepta 183 2.4% (1.9–2.9%)	hepta 183 6.0% (4.0–7.9%)	hepta 182 3.1% (0.9–5.3%)	hexa 153 5.2% (3.7–7.6%)	penta 99 (DNQ)	tetra 49 0.8% (0.7–1.0%)	di 4 28% (0–55%)
	hepta 176 0.9% (0.7–1.2%)		hepta 180 1.2% (0.4–2%)	hexa 149 4.3% (3.4–6.1%)	tetra 47 (DNQ)	tetra 48 18% (13–27%)	
			hepta 175* 2.1% (1.1–3.1%)	hexa 144 0.7% (0.6–0.7%)		tetra 47 (DNQ)	
				penta 99 (DNQ)		tri 18 (DNQ)	
<i>D. restrictus</i>	hepta 184 4.6% (2.4–6.7%)	hepta 187* 7.6% (7.1–8.1%)	hepta 191 4.1% (1.0–6.0%)	hexa 154 1.5% (0.7–2.9%)	penta 101 2.9% (0.7–2.9%)	tetra 49 2.5% (1.3–3.7%)	tri 17 2.8% (2.6–2.9%)
	hepta 183 2.4% (2.2–2.6%)	hepta 183 6.2% (6.0–6.5%)	hepta 183 7.5% (2.2–11.8%)	hexa 153 0.6% (0.2–1.4%)	penta 99 7.0% (0.2–1.4%)	tetra 48 22% (17–26%)	di 4 84% (72–96%)
	hepta 176 2.6% (1.1–3.4%)	hepta 180 0.2% (0.24–0.25%)	hepta 182 2.3% (0.8–3.2%)	hexa 149 1.9% (0.2–5.1%)	tetra 48 3.2%	tetra 47 1.1% (0.8–1.4%)	
			hepta 175* 14.2% (4.2–22%)	hexa 144 0.6% (0.2–1.4%)			
				hexa 139 0.2% (0.1–0.3%)			

^a The products from each substrate are listed in separate columns beneath the substrate name. The products are divided by culture, listed on the left. The average mole percent of each product at the end of the sampling period is listed next to the name. The range of mole percents for the replicate samples where available is listed in parentheses beneath the average value. The *D. restrictus* 47 data were collected after two weeks and the ANAS195 197 data were collected after 6 months. DNQ indicates that the congener was detected but was not quantifiable. * Indicates congener that is presumptively identified due to lack of available standards.

the extent of debromination varied greatly between cultures and between congeners (Table 1). However, the debromination pathway for each congener was surprisingly similar for all tested cultures. For simplification in describing the pathways, we assumed that there was no isomeric rearrangement of bromines around the PBDE ring during debromination. Except for two unavailable congeners, all substrate and product PBDE peaks were matched with standards for specific identification and quantification. Figure 3 summarizes the observed debromination pathway for all tested congeners. Based on the geometries of the substrate molecules, possible debromination products ranged from two to eight possible congeners, although in most cases many of these congeners were never produced. All cultures exhibited preferences for removing bromines at certain positions, typically the meta and para bromines.

In some cases, due to the synthesis process, the substrates contained impurities that were also possible debromination products. For example, octa 203 contained trace (around 0.1 mol percent) quantities of heptas 183 and 187 and hepta 183 contained hexa 153 and 154. However, in each case we were able to confirm that these PBDEs were being biologically produced because the concentrations of the products increased significantly (at least 2-fold) in the active bottles while remaining constant in control bottles during the three month incubation.

Biotransformation of the octa congeners was rather slow and limited for all tested cultures, in most cases with less

than 10 mol% of the original substrate being degraded within three months. Octa 197 was debrominated by ortho substitution to hepta 183, by meta substitution to hepta 184, and by para substitution to 176 by all three cultures. All congeners were generated in approximately equal quantities, cumulatively amounting to 9 mol% or less of the total recovered molar concentration of PBDEs at the end of three months. Octa 203 debromination produced two major congeners for all three cultures: hepta 183 and presumptively identified hepta 187—the meta and para bromine removal products respectively. Both peaks represented between 2 and 7 mol% after three months. An ortho debromination product, hepta 180, was produced in trace quantities, accounting for only 0.3 mol%. Congener 187 is considered presumptively identified because that congener was not available for purchase. However, given the geometry of the bromines positioned on the octa 203 molecule, there are only six possible debromination products (heptas 180, 183, 187, 190, 185, and 181) and the availability of standards for all of these except for 187 facilitated identification by elimination. That is, of the three product peaks detected in all samples, two have matching standards (180 and 183), while the other three standards (191, 185, and 181) did not match the third product peak, leaving 187 as the lone possible candidate for the unidentified peak.

Four congeners were identified as debromination products from octa 196, including heptas 183, 182, and presumptively identified 175 for all species. *D. restrictus* also




FIGURE 3. Summarized PBDE debromination pathway for all tested congeners. Highlighted molecules are those that were applied as initial substrate. The cultures that produced each congener are listed by the reaction arrows. Note that since the rings can rotate around the oxygen bond, some congener depictions are obvious with some pathways (203 to 183) and not with others (197 to 183). Note also that molecules have sometimes been flipped horizontally in the diagram (e.g., 196 to 183). Asterisk (*) indicates congener that is presumptively identified due to lack of available standards.

produced the ortho debromination product 191 whereas *D. hafniense* and ANAS195 produced 180—a different ortho product. One of the possible debromination products of octa 196, hepta 174 was not available for purchase and therefore the production of this congener could not be verified. Although one of the product peaks perfectly matches the 175 standard, it could not be excluded that this peak might be 174. According to relative retention time estimates by Wang et al. (25) and Rayne and Ikononou (26), 175 and 174 have very similar retention times and might coelute even on a two-dimensional GC. Furthermore, both 175 and 174 are para bromine removal products and thus equally likely possibilities. However, 175 is a doubly flanked para removal whereas 174 is a singly flanked para removal. Since doubly flanked bromines are preferred over singly flanked ones in the other PBDE pathways, we have presumptively designated hepta 175 as the product.

Hepta 183 was debrominated relatively rapidly by all three cultures, producing many products including hexas 154 (meta removal), 153 (ortho removal), 149, and 144 (para removal products). Additionally, 183 was debrominated to the meta substitution product hexa 139 by only *D. restrictus*. Oddly,

the two para removal products varied significantly in concentrations, with 149 being a major product with concentrations up to 4 mol% and 144 produced only in minor quantities, representing on average 0.5 mol%. Hexa 153 was debrominated to pentas 99 and 101 by all three cultures. Penta 118 required an ortho bromine removal and was produced only by the ANAS195 culture in trace quantities at around 0.6 mol%.

In all cultures, penta 99 was debrominated to tetras 47 and 49 by meta and para substitution respectively. ANAS195 and *D. hafniense* also produced tetra 66, an ortho bromine removal product, whereas *D. restrictus* and *D. hafniense* substituted a para bromine, producing tetra 48. Tetras 49 and 48 were the predominant congeners, representing up to 22 mol%, whereas 47 and 66 were detected at concentrations less than 1 mol% after three months. In particular, tetra 47 appeared only at trace concentrations probably because it is so biodegradable. When tetra 47 was exposed to bacteria as a substrate, it was biotransformed relatively rapidly particularly in the *D. hafniense* and *D. restrictus* cultures where the original congener was almost completely debrominated by the conclusion of the experiment. *D. hafniense*

and *D. restrictus* produced only tri-BDE 17, which was quickly debrominated further to di-BDE 4. This di-BDE congener represented a major product at 28 and 83 mol% in the *D. hafniense* and *D. restrictus* cultures, respectively. It is possible that further debromination products such as mono-BDEs were produced in the *D. hafniense* and *D. restrictus* cultures, but these congeners were never detected, perhaps due to the low sensitivity of electron capture detection to mono-BDEs, interference with compounds in the bacterial medium that elute at the same time as mono-BDEs, and low mono-BDE concentrations. The ANAS195 culture, in addition to the para substitution product tri-BDE 17, generated the ortho bromine removal product tri-BDE 28, although in trace concentrations (0.2 mol%). Di-BDE 4 was never observed in the ANAS195 samples.

Although Figure 3 depicts only the products of one bromine removal for each tested congener, further debromination products were frequently detected. These additional products are not represented in the pathway schematic as it is not always evident which congener they were derived from, since often multiple debromination pathways (para, meta, ortho) are possible. These subsequent debromination products are listed in Table 1 along with all of the single bromine removal products for each congener and bacterial culture.

Growth of the bacteria and degradation of their primary chlorinated electron acceptor were monitored during all experiments. *D. restrictus* and *D. hafniense* degraded their growth substrates within one month, whereas PCE dechlorination by ANAS195 continued throughout the entire three month incubation. The *D. restrictus* and *D. hafniense* cultures started at cell densities of 10^6 and 10^5 cells/mL, respectively, and reached average maximum cell densities of 10^7 cells/mL within one month and then began to decrease. The *Dehalococcoides* strains within the ANAS195 culture started at cell densities of 10^6 cells/mL and reached 10^7 cells/mL after two months and remained constant thereafter. In all experiments and all cultures, PBDE debromination continued throughout the three month period.

To determine whether chlorinated substrates were required to induce PBDE debromination, additional experiments were conducted with these three cultures exposed to the octa-BDE mixture in the absence of PCE or pentachlorophenol. No debromination was observed in the *D. restrictus* and *D. hafniense* samples, whereas debromination was observed in the ANAS195 culture.

Discussion

All of the PBDE congeners evaluated in this study were found to be debrominated to some degree by anaerobic dehalogenating microorganisms. The debromination pathways were found to be generally consistent across the different cultures, with exceptions typically represented by a single congener produced in trace quantities. In general, debromination was slower for the more brominated congeners, such as the octas, than for the less brominated penta and tetras—a trend that has also been observed with PCBs (27) and that can be explained by the increasing hydrophobicity of PBDEs as the number of bromines increases. The *D. hafniense* and *D. restrictus* cultures were able to dehalogenate almost all of the tetra 47 they were exposed to (the least hydrophobic PBDE tested), while transforming typically less than 10 mol% of the octa congeners in the same time period.

Para and meta bromines were preferentially removed from the rings by the cultures, often with multiple para or meta bromine removal products being formed. The most commonly substituted bromines were those that are double flanked. In the case of octa-BDE 203, the doubly flanked meta and para bromines were removed whereas the two

other singly flanked para and meta bromines were never removed. Given that repulsion between adjacent bromines results in increased enthalpies of formation, it is not surprising that doubly flanked bromines are removed more frequently than singly flanked bromines (28).

Ortho bromines were also frequently removed. Typically, however, the ortho substitution products were minor ones compared to the para and meta bromine removal products in terms of concentration and were sometimes not even produced (as is evident with the lack of production of tri 28 from tetra 47 and penta 118 from hexa 153 for both the *D. hafniense* and *D. restrictus* cultures). Furthermore, as is evident with hepta 183, only one of the ortho-bromine removal products (hexa 153) was formed, while all the possible meta and para bromines were removed. This emphasizes the difficulty in removing ortho halogens, as had been observed with PCBs and polybrominated biphenyls (PBBs) (29–31). The most notable pathway differences between species occurred with ortho bromine removals. For example, *D. hafniense* and ANAS195 preferred to remove the ortho bromine on the fully brominated ring of octa 196, creating hepta 180 whereas *D. restrictus* preferred the ortho removal on the other ring, creating hepta 191. The ANAS195 culture was better than *D. hafniense* and *D. restrictus* at ortho bromine removals, which might be explained by the fact that *D. ethenogenes* 195 is known to dechlorinate chlorophenols by ortho removal in the presence of an adjacent meta chlorine (32). Oddly, the PBDE dehalogenation patterns for *D. hafniense* differed from its chlorophenol dehalogenation patterns in that it prefers to remove the ortho chlorines from chlorophenols, whereas it preferentially removes meta and para bromines from PBDEs (16).

Our study is the first to show that bacteria in the *Desulfitobacterium* and *Dehalobacter* genera can transform PBDEs. In particular *Dehalobacter* had not been previously shown to dehalogenate aromatic compounds, unlike *Dehalococcoides* and *Desulfitobacterium* (33–35). Both *D. restrictus* and *D. hafniense* exhibited more extensive PBDE debromination producing higher concentrations of products compared to the ANAS195 culture, which degraded all congeners much more slowly. This difference is perhaps due to the faster growth exhibited in the *D. restrictus* and *D. hafniense* cultures. *D. restrictus* and *D. hafniense* were both capable of rapid removal of multiple bromines as is particularly evident with tetra 47, which both cultures transformed to mostly di-BDE 4 within two weeks in the case of *D. restrictus* and within three months in the case of *D. hafniense*.

The debromination results using individual congeners described here correlate well with previously reported debromination of an octa mixture in that many of the same products were generated (8). For example, the dominant hexas generated from hepta 183 in this study (hexas 154, 153, 149, and 144) were the same as those generated from debromination of the octa-BDE mixture. The observed debromination products of hexa 153 (pentas 99 and 101) were also detected in that study, as were two of the debromination products of penta 99 (tetras 47 and 49) and the tetra 47 debromination product tri 17. There were also other products in the octa mixture samples that did not match our individual congener studies. This was expected given that there were additional minor components in the octa mixture which were not evaluated in this study. Furthermore, since the octa mixture samples were incubated for almost a year, many of the generated products likely resulted from multiple bromine removal steps.

An interesting aspect of these results is their striking similarity to PCB and PBB dehalogenation studies. The prevalent removal strategies observed in this study were meta and para debrominations, as is common with PCBs and PBBs

(29–31). Specifically, Kuipers et al. found that octa-CB 203 produced hepta-CB 183 similar to what was shown here for PBDEs (36). Hepta-CB 181 was also generated, whereas this congener was never detected in PBDE samples. Octa-CB 196 was dechlorinated to hepta-CB 183, as was observed with the PBDE analog. Morris et al. found that with PBBs anaerobic sediment microorganisms debrominated hexa-BB 153 to tetra-BBs 49 and 47 and tetra-BB 47 to tri-BB 17, exactly as has been observed in this study (30).

Dehalococcoides is known to dechlorinate PCBs (27, 34, 37). Fagervold et al. found that microbial microcosms dechlorinated hepta-CB 183 to hexa-CB 154, hexa-CB 153 to penta-CB 99, and penta-CB 99 to tetra-CB 47, as analogous to the observation in our study (27). One notable difference between PCB and PBDE degradation, however, is that ortho dechlorination appears to be quite rare with PCBs, whereas ortho debromination products were detected in all of our experiments.

The extent of PBDE dehalogenation observed in this study, in which nM quantities were transformed over the course of months, is orders of magnitude less than reported PCB dechlorination, in which μ M quantities were transformed over the same time period (27, 33, 34). This may in part be due to higher cell densities of dechlorinating organisms, which ranged from 10^8 to 10^9 cells/mL in pure and mixed cultures. However, it may also be due to the approximately 10-fold higher hydrophobicity of PBDEs compared to PCBs (9, 38).

As evidenced by the lack of debromination observed with *D. tiedjei* DCB-1, not all dehalogenating bacteria are capable of catalyzing PBDE debromination. Although we have not identified the specific enzymes responsible for these reactions, it is possible that the reductive dehalogenases responsible for PCE and pentachlorophenol degradation are involved in debromination because experiments with *D. restrictus* and *D. hafniense* in which PCE or pentachlorophenol were not added as electron acceptor generated no detectable PBDE debromination activity. These results suggest that either the debrominating enzymes were not induced by the PBDEs alone or that the PBDE transformation by these isolates is cometabolic, requiring the concomitant presence of energy-generating electron acceptors. In contrast, the ANAS195 culture was able to debrominate PBDEs in the absence of PCE, although the mechanism behind this reaction is unclear.

In conclusion, the similarity of debromination pathways exhibited by three very different bacterial cultures analyzed in this study suggests that pathways for PBDE debromination are conserved across some species and genera. An additional notable observation is that although tetra 47 and penta 99, two of the most toxic PBDE congeners, are products of higher congeners, they are also relatively readily debrominated by anaerobic bacteria to at least di-BDEs. This suggests that it is possible that their accumulation could be transient, leading to the generation of lesser toxic final products.

Acknowledgments

We thank Patrick Lee for providing the ANAS culture and Professor Steve Zinder for providing *D. ethenogenes* 195. Funding was provided by the University of California Center for Water Resources and the NIEHS Superfund Basic Research Program under grant ES04705. K.R. also thanks Dr. Gregory Cost for excellent scientific discussions and advice.

Literature Cited

- (1) Alaei, M.; Arias, P.; Sjödin, A.; Bergman, A. An overview of commercially used brominated flame retardants, their applications, their use patterns in different countries/regions and possible modes of release. *Environ. Int.* **2003**, *29* (6), 683–689.
- (2) Wit, D. An overview of brominated flame retardants in the environment. *Chemosphere* **2002**, *46*, 583–674.
- (3) Zhou, T.; Taylor, M. M.; DeVito, M. J.; Crofton, K. A., Jr. Developmental exposure to brominated diphenyl ethers results in thyroid hormone disruption. *Toxicol. Sci.* **2002**, *66* (1), 105–116.
- (4) Norén, K.; Meironyté, D. Certain organochlorine and organobromine contaminants in Swedish human milk in perspective of past 20–30 years. *Chemosphere* **2000**, *40* (9–11), 1111–1123.
- (5) Oros, D. R.; Hoover, D.; Rodigari, F.; Crane, D.; Sericano, J. Levels and distribution of polybrominated diphenyl ethers in water, surface sediments, and bivalves from the San Francisco Estuary. *Environ. Sci. Technol.* **2005**, *39* (1), 33–41.
- (6) Forum, B. S. a. E. EU chemicals assessment authorities confirm no restrictions on Deca-BDE and support ongoing science. <http://bsef.com/newsmanager/newstemplate.php?id=272>, December 17, 2007.
- (7) Gerecke, A. C.; Hartmann, P. C.; Heeb, N. V.; Kohler, H. P.; Giger, W.; Schmid, P.; Zennegg, M.; Kohler, M. Anaerobic degradation of decabromodiphenyl ether. *Environ. Sci. Technol.* **2005**, *39* (4), 1078–1083.
- (8) He, J.; Robrock, K. R.; Alvarez-Cohen, L. Microbial reductive debromination of polybrominated diphenyl ethers (PBDEs). *Environ. Sci. Technol.* **2006**, *40* (14), 4429–4434.
- (9) Darnerud, P. O.; Eriksen, G. S.; Johannesson, T.; Larsen, P. B.; Viluksela, M. Polybrominated diphenyl ethers: Occurrence, dietary exposure, and toxicology. *Environ. Health Perspect.* **2001**, *109*, 49–68.
- (10) Korytár, P.; Covaci, A.; Leonards, P. E. G.; de Boer, J.; Brinkman, U. A. T. Comprehensive two-dimensional gas chromatography of polybrominated diphenyl ethers. *J. Chromatogr. A* **2005**, *1100* (2), 200–207.
- (11) La Guardia, M. J.; Hale, R. C.; Harvey, E. Detailed polybrominated diphenyl ethers (PBDE) congener composition of the widely used penta-, octa- and deca-PBDE technical flame-retardant mixtures. *Environ. Sci. Technol.* **2006**, *40* (20), 6247–6254.
- (12) Richardson, R. E.; Bhupathiraju, V. K.; Song, D. L.; Goulet, T. A.; Alvarez-Cohen, L. Phylogenetic characterization of microbial communities that reductively dechlorinate TCE based upon a combination of molecular techniques. *Environ. Sci. Technol.* **2002**, *36* (12), 2652–2662.
- (13) Holmes, V. F.; He, J. Z.; Lee, P. K. H.; Alvarez-Cohen, L. Discrimination of multiple *Dehalococcoides* strains in a trichloroethene enrichment by quantification of their reductive dehalogenase genes. *Appl. Environ. Microbiol.* **2006**, *72* (9), 5877–5883.
- (14) Maymó-Gatell, Chien, Y. T.; Gossett, J. M.; Zinder, S. H. Isolation of a bacterium that reductively dechlorinates tetrachloroethene to ethene. *Science* **1997**, *276* (5318), 1568–1571.
- (15) Holliger, C.; Hahn, D.; Harmsen, H.; Ludwig, W.; Schumacher, W.; Tindall, B.; Vazquez, F.; Weiss, N.; Zehnder, A. J. B. *Dehalobacter restrictus* gen. nov. and sp. nov., a strictly anaerobic bacterium that reductively dechlorinates tetra- and trichloroethene in an anaerobic respiration. *Arch. Microbiol.* **1998**, *169* (4), 313–321.
- (16) Bouchard, B.; Beaudet, R.; Villemur, R.; McSweeney, G.; Lepine, F.; Bisailon, J. G. Isolation and characterization of *Desulfotobacterium frappieri* sp. nov., an anaerobic bacterium which reductively dechlorinates pentachlorophenol to 3-chlorophenol. *Int. J. Syst. Bacteriol.* **1996**, *46* (4), 1010–1015.
- (17) Sanford, R. A.; Cole, J. R.; Löffler, F. E.; Tiedje, J. M. Characterization of *Desulfotobacterium chlororespirans* sp. nov., which grows by coupling the oxidation of lactate to the reductive dechlorination of 3-chloro-4-hydroxybenzoate. *Appl. Environ. Microbiol.* **1996**, *62* (10), 3800–3808.
- (18) Utkin, I.; Woese, C.; Wiegand, J. Isolation and characterization of *Desulfotobacterium dehalogenans* gen. nov., sp. nov., an anaerobic bacterium which reductively dechlorinates chlorophenolic compounds. *Int. J. Syst. Bacteriol.* **1994**, *44* (4), 612–619.
- (19) Deweerdt, K. A.; Mandelco, L.; Tanner, R. S.; Woese, C. R.; Sufita, J. M. *Desulfomonile tiedjei* gen. nov. and sp. nov., a Novel Anaerobic, Dehalogenating, Sulfate-Reducing Bacterium. *Arch. Microbiol.* **1990**, *154* (1), 23–30.
- (20) He, J. Z.; Ritalahti, K. M.; Aiello, M. R.; Löffler, F. E. Complete detoxification of vinyl chloride by an anaerobic enrichment culture and identification of the reductively dechlorinating population as *Dehalococcoides* species. *Appl. Environ. Microbiol.* **2003**, *69* (2), 996–1003.
- (21) Freeborn, R. A.; West, K. A.; Bhupathiraju, V. K.; Chauhan, S.; Rahm, B. G.; Richardson, R. E.; Alvarez-Cohen, L. Phylogenetic

- analysis of TCE-dechlorinating consortia enriched on a variety of electron donors. *Environ. Sci. Technol.* **2005**, 39 (21), 8358–8368.
- (22) Levesque, M.-J.; La Boissiere, S.; Thomas, J.-C.; Beaudet, R.; Villemur, R. Rapid method for detecting *Desulfotobacterium frappieri* PCP-1 in soil by polymerase chain reaction. *Appl. Microbiol. Biotechnol.* **1997**, 47, 719–725.
 - (23) Staskal, D. F.; Diliberto, J. J.; DeVito, M. J.; Birnbaum, L. S. Toxicokinetics of BDE 47 in female mice: Effect of dose, route of exposure, and time. *Toxicol. Sci.* **2005**, 83 (2), 215–223.
 - (24) Eriksson, P.; Viberg, H.; Jakobsson, E.; Örn, U.; Fredriksson, A. A brominated flame retardant, 2,2',4,4',5-pentabromodiphenyl ether: Uptake, retention, and induction of neurobehavioral alterations in mice during a critical phase of neonatal brain development. *Toxicol. Sci.* **2002**, 67 (1), 98–103.
 - (25) Wang, Y.; Li, A.; Liu, H.; Zhang, Q.; Ma, W.; Song, W.; Jiang, G. Development of quantitative structure gas chromatographic relative retention time models of seven stationary phases for 209 polybrominated diphenyl ether congeners. *J. Chromatogr. A* **2006**, 1103, 314–328.
 - (26) Rayne, S.; Ikononou, M. G. Predicting gas chromatographic retention times for the 209 polybrominated diphenyl ether congeners. *J. Chromatogr. A* **2003**, 1016 (2), 235–248.
 - (27) Fagervold, S. K.; May, H. D.; Sowers, K. R. Microbial reductive dechlorination of aroclor 1260 in Baltimore Harbor sediment microcosms is catalyzed by three phylotypes within the phylum Chloroflexi. *Appl. Environ. Microbiol.* **2007**, 73 (9), 3009–3018.
 - (28) Zeng, X.; Freeman, P. K.; Vasil'ev, Y. V.; Voinov, V. G.; Simonich, S. L.; Barofsky, D. F. Theoretical calculation of thermodynamic properties of polybrominated diphenyl ethers. *J. Chem. Eng. Data* **2005**, 50 (5), 1548–1556.
 - (29) Bedard, D. L.; Van Dort, H. M. Complete reductive dehalogenation of brominated biphenyls by anaerobic microorganisms in sediment. *Appl. Environ. Microbiol.* **1998**, 64 (3), 940–947.
 - (30) Morris, P.; Quensen, J. F.; Tiedje, J. M.; Boyd, S. A. Reductive debromination of the commercial polybrominated biphenyl mixture Firemaster BP6 by aerobic microorganisms from sediments. *Appl. Environ. Microbiol.* **1992**, 58 (10), 3249–3256.
 - (31) Wiegel, J.; Wu, Q. Z. Microbial reductive dehalogenation of polychlorinated biphenyls. *FEMS Microbiol. Ecol.* **2000**, 32 (1), 1–15.
 - (32) Adrian, L.; Hansen, S. K.; Görisch, H.; Zinder, S. H. Growth of *Dehalococcoides* strains with chlorophenols as electron acceptors. *Environ. Sci. Technol.* **2007**, 41 (7), 2318–2323.
 - (33) Bedard, D. L.; Ritalahti, K. A.; Löffler, F. E. The *Dehalococcoides* population in sediment-free mixed cultures metabolically dechlorinates the commercial polychlorinated biphenyl mixture aroclor 1260. *Appl. Environ. Microbiol.* **2007**, 73 (8), 2513–2521.
 - (34) Fennell, D. E.; Nijenhuis, I.; Wilson, S. F.; Zinder, S. H.; Häggblom, M. M. *Dehalococcoides ethenogenes* strain 195 reductively dechlorinates diverse chlorinated aromatic pollutants. *Environ. Sci. Technol.* **2004**, 38 (7), 2075–2081.
 - (35) Wiegel, J.; Zhang, X. M.; Wu, Q. Z. Anaerobic dehalogenation of hydroxylated polychlorinated biphenyls by *Desulfotobacterium dehalogenans*. *Appl. Environ. Microbiol.* **1999**, 65 (5), 2217–2221.
 - (36) Kuipers, B.; Cullen, W. R.; Mohn, W. W. Reductive dechlorination of nonachlorobiphenyls and selected octachlorobiphenyls by microbial enrichment cultures. *Environ. Sci. Technol.* **1999**, 33 (20), 3579–3585.
 - (37) Bedard, D. L.; Bailey, J. J.; Reiss, B. L.; Jerzak, G. V. Development and characterization of stable sediment-free anaerobic bacterial enrichment cultures that dechlorinate Aroclor 1260. *Appl. Environ. Microbiol.* **2006**, 72 (4), 2460–2470.
 - (38) Ballschmiter, K.; Klingler, D.; Ellinger, S.; Hackenberg, R. High-resolution chromatography retention data as a basis for estimation of the octanol-water distribution coefficients (K_{ow}) of PCB: the effect of experimental conditions. *Anal. Bioanal. Chem.* **2005**, 382 (8), 1859–1870.

ES0720917